

The Stochastic Zakharov system in dimension 1

Grégoire Barrué,
ENS Rennes,

Email : gregoire.barrue@ens-rennes.fr

Mots Clés : Équations aux dérivées partielles stochastiques, système de Zakharov, équation de Schrödinger, processus de Wiener ...

Biographie – J'ai fait mes études à l'ENS Rennes et y suis maintenant en thèse sous la direction d'Arnaud Debussche et Anne De Bouard. Ma thèse s'inscrit dans l'étude d'équations aux dérivées partielles stochastiques dans un contexte d'approximation diffusion, et plus particulièrement du système de Zakharov et de l'équation de Schrödinger.

Resumé :

The Zakharov system is a simplified model for the description of Langmuir oscillations in a ionized plasma (for example the ionosphere). Langmuir waves are rapid oscillations of the electron density.

We are interested in the stochastic Zakharov system, with a damping term :

$$\begin{cases} i\partial_t u = -\partial_x^2 u + nu \\ \varepsilon^2 \partial_t^2 n + \alpha \varepsilon \partial_t n = \partial_x^2(n + |u|^2) + \phi \frac{dW(t)}{dt} \end{cases} \quad (1)$$

where ϕ is a regularizing operator, and $\frac{dW(t)}{dt}$ is a space-time white noise. The function u represents the envelope of the electric field, and n denotes the deviation of the ion density from its mean mode. The term ε^2 refers to the inverse of the ion speed. The stochastic perturbation of this system can be understood as random external fluctuations, as for instance thermal fluctuations.

The deterministic version of this system (without damping term) was studied first by Sulem and Sulem in [10] where they proved global well-posedness in dimension 1 for regular initial data. More refinements were made by Added and Added in [1] to get global well-posedness in dimension 2 for small initial data. We can also cite [6] by Ginibre, Tsutsumi and Velo which proved local well-posedness in the energy space. The limit $\varepsilon \rightarrow 0$ of this system is studied for example in [2] where Added and Added showed the convergence to a cubic Nonlinear Schrödinger equation. In [8], Masmoudi and Nakanishi have provided a simpler proof of this convergence. We adapted the work of [10] to show almost sure global well-posedness in dimension 1, and we added a damping term to prove that for regular initial data, the solution u^ε of the stochastic Zakharov system converges in law to a stochastic Nonlinear Schrödinger equation :

$$i\partial_t u = -\partial_x^2 u - |u|^2 u + u\psi \circ \frac{dW(t)}{dt}. \quad (2)$$

To study this stochastic problem, we have to place ourselves in the context of approximation-diffusion, because the process (n^ε) has a non-zero time correlation length and we want it to converge to a white noise. The main difficulty comes from the fact that the energy of our system

$$H(u, n) = \|\partial_x u\|_{L^2}^2 + \frac{1}{2} (\|n\|_{L^2}^2 + \|\varepsilon \partial_t \partial_x^{-1} n\|_{L^2}^2) + \int_{\mathbb{R}} n|u|^2 dx \quad (3)$$

is no longer preserved, and more importantly it has a singular evolution as ε goes to 0. Thus we can not just adapt the work in [2], and we need to rescale our system to use the Perturbed Test Function method in order to correct the infinitesimal generator of $(u^\varepsilon, n^\varepsilon)$ and pass to the limit in the martingale problem.

Références

- [1] Hélène Added and Stéphane Added. Existence globale de solutions fortes pour les équations de la turbulence de Langmuir en dimension 2. *C. R. Acad. Sci. Paris Sér. I Math.*, 299(12):551–554, 1984.
- [2] Hélène Added and Stéphane Added. Equations of Langmuir turbulence and nonlinear Schrödinger equation: smoothness and approximation. *J. Funct. Anal.*, 79(1):183–210, 1988.
- [3] P. Billingsley. *Convergence of probability measures*. John Wiley Sons, 1999.
- [4] Giuseppe Da Prato and Jerzy Zabczyk. *Stochastic equations in infinite dimensions*, volume 152 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition, 2014.
- [5] A. de Bouard and A. Debussche. The stochastic nonlinear Schrödinger equation in H^1 . *Stochastic Anal. Appl.*, 21(1):97–126, 2003.
- [6] J. Ginibre, Y. Tsutsumi, and G. Velo. On the Cauchy problem for the Zakharov system. *J. Funct. Anal.*, 151(2):384–436, 1997.
- [7] Tosio Kato. On nonlinear Schrödinger equations. *Ann. Inst. H. Poincaré Phys. Théor.*, 46(1):113–129, 1987.
- [8] Nader Masmoudi and Kenji Nakanishi. Energy convergence for singular limits of Zakharov type systems. *Invent. Math.*, 172(3):535–583, 2008.
- [9] P. Nom1, P. Nom2, and P. Nom3. Le titre de l'article. *Journal mathématique de quelque chose*, 1(1), 2020.
- [10] Catherine Sulem and Pierre-Louis Sulem. Quelques résultats de régularité pour les équations de la turbulence de Langmuir. *C. R. Acad. Sci. Paris Sér. A-B*, 289(3):A173–A176, 1979.